

Casa Grande drug use data

Based on Arizona Youth Survey, Arizona Criminal Justice Commission

Note: an 'x' indicates that question was not asked that year

30 day drug use

Substance	Question	grade 10			grade 12		
		2004	2008		2004	2006	
		CG	CG	AZ	CG	CG	AZ
Alcohol	had alcoholic beverages (beer, wine or hard liquor) to drink - more than just a few sips?	41.3	41.7	37.7	46	50	47
Cigarettes	smoked cigarettes?	18.1	16.8	16.6	23.1	23.8	21.8
chewing tobacco	used smokeless tobacco (chew, snuff, plug, dipping tobacco, chewing tobacco?)	2.8	4.3	4.6	5.5	4.8	5.4
Marijuana	used marijuana?	17.1	16	15.1	23.7	17.1	18.1
Hallucinogens	used LSD or other hallucinogens?	2.2	3	1.9	1.7	5.1	1.7
Cocaine	used cocaine or crack?	5	4.7	2.2	0	7.3	3.3
Inhalants	sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high?	5.6	3.5	3	0	2.4	1.7
Stimulants & Meth	used amphetamines (meth, crystal, crank) without a doctor telling you to take them?	4.5	0	0.6	6.1	2.4	2.6
Heroin/Other opiates	used heroin or other opiates?	1.7	2.1	0.07	0	0	0.8
Ecstasy	used Ecstasy (X, E, or MDMA)?	1.7	1.8	1.5	0	0	0.9
Steroid	used steroids or anabolic steroids	x	1.2	1	0	0	1
Prescription Pain Relievers	used prescription pain relievers (such as Vicodin, OxyContin, Percocet or Codeine) w/o doctor telling you to take them?	x	13	9.4	x	x	x
Prescription stimulants	used prescription stimulants (such as Ritalin, Adderall, or Dexedrine) without a doctor telling you to take them?	x	3.6	2.9	x	2.4	2.6
Prescription sedatives	used prescription sedatives (tranquilizers, such as Valium or Xanax, barbiturates or sleeping pills) w/o a doctor telling you to take them?	8.4	3.3	4.5	8.8	4.9	x
Prescription drugs	combined stimulants, sedative and pain reliever	x	14.5	12.2	x	5	8.1
Over the counter	used over the counter drugs such as cough syrup, cold medicine, or diet pills for the purposes of getting high?	x	10.1	6.4	x	x	x